

Anisotropic cosmological models with a perfect fluid and a Λ term

Bijan Saha

*Laboratory of Information Technologies
Joint Institute for Nuclear Research, Dubna
141980 Dubna, Moscow region, Russia**

(Dated: February 7, 2008)

We consider a self-consistent system of Bianchi type-I (BI) gravitational field and a binary mixture of perfect fluid and dark energy given by a cosmological constant. The perfect fluid is chosen to be the one obeying either the usual equation of state, i.e., $p = \zeta\varepsilon$, with $\zeta \in [0, 1]$ or a van der Waals equation of state. Role of the Λ term in the evolution of the BI Universe has been studied.

PACS numbers: 04.20.Ha, 03.65.Pm, 04.20.Jb

Keywords: Bianchi type I (BI) model, perfect fluid, van der Waals fluid

I. INTRODUCTION

In view of its importance in explaining the observational cosmology many authors have considered cosmological models with dark energy. In a recent paper Kremer [1] has modelled the Universe as a binary mixture whose constituents are described by a van der Waals fluid and by a dark energy density. Zlatev *et al.* [2] showed that "tracker field", a form of quintessence, may explain the coincidence, adding new motivation for the quintessence scenario. The fate of density perturbations in a Universe dominated by the Chaplygin gas, which exhibit negative pressure was studied by Fabris *et al.* [3]. Model with Chaplygin gas was also studied in the Refs. [4, 5]. In doing so the author considered a spatially flat, homogeneous and isotropic Universe described by a Friedmann-Robertson-Walker (FRW) metric. Since the theoretical arguments and recent experimental data support the existence of an anisotropic phase that approaches an isotropic one, it makes sense to consider the models of Universe with anisotropic back-ground in presence of dark energy. The simplest of anisotropic models, which nevertheless rather completely describe the anisotropic effects, are Bianchi type-I (BI) homogeneous models whose spatial sections are flat but the expansion or contraction rate is direction-dependent. In a number of papers, e.g., [6, 7], we have studied the role of a Λ term in the evolution of a BI space-time in presence of spinor and/or scalar field with a perfect fluid satisfying the equation of state $p = \zeta\varepsilon$. In this paper we study the evolution of an initially anisotropic Universe given by a BI spacetime in presence of a perfect fluid obeying not only $p = \zeta\varepsilon$, but also the Van der Waals equation of state.

*Electronic address: saha@thsun1.jinr.ru, bijan@jinr.ru; URL: <http://thsun1.jinr.ru/~saha/>

II. BASIC EQUATIONS

The Einstein field equation on account of the cosmological constant we write in the form

$$R^\nu_\mu - \frac{1}{2}\delta^\nu_\mu R = \kappa T^\nu_\mu + \delta^\nu_\mu \Lambda. \quad (2.1)$$

Here R^ν_μ is the Ricci tensor, R is the Ricci scalar and κ is the Einstein gravitational constant. As was mentioned earlier, Λ is the cosmological constant. To allow a steady state cosmological solution to the gravitational field equations Einstein [8] introduced a fundamental constant, known as cosmological constant or Λ term, into the system. Soon after E. Hubble had experimentally established that the Universe is expanding, Einstein returned to the original form of his equations citing his temporary modification of them as the biggest blunder of his life. Λ term made a temporary comeback in the late 60's. Finally after the pioneer paper by A. Guth [9] on inflationary cosmology researchers began to study the models with Λ term with growing interest. Note that in our previous papers [6, 7] we studied the Einstein field equations where the cosmological term appears with a negative sign. Here following the original paper by Einstein and one by Sahni [10] we choose the sign to be positive. In this paper a positive Λ corresponds to the universal repulsive force, while a negative one gives an additional gravitational force. Note that a positive Λ is often taken to a form of dark energy.

We study the gravitational field given by an anisotropic Bianchi type I (BI) cosmological model and choose it in the form:

$$ds^2 = dt^2 - a^2 dx^2 - b^2 dy^2 - c^2 dz^2, \quad (2.2)$$

with the metric functions a, b, c being the functions of time t only.

The Einstein field equations (2.1) for the BI space-time in presence of the Λ term now we write in the form

$$\frac{\ddot{b}}{b} + \frac{\ddot{c}}{c} + \frac{\dot{b}\dot{c}}{b c} = \kappa T_1^1 + \Lambda, \quad (2.3a)$$

$$\frac{\ddot{c}}{c} + \frac{\ddot{a}}{a} + \frac{\dot{c}\dot{a}}{c a} = \kappa T_2^2 + \Lambda, \quad (2.3b)$$

$$\frac{\ddot{a}}{a} + \frac{\ddot{b}}{b} + \frac{\dot{a}\dot{b}}{a b} = \kappa T_3^3 + \Lambda, \quad (2.3c)$$

$$\frac{\dot{a}\dot{b}}{a b} + \frac{\dot{b}\dot{c}}{b c} + \frac{\dot{c}\dot{a}}{c a} = \kappa T_0^0 + \Lambda. \quad (2.3d)$$

Here over-dot means differentiation with respect to t . The energy-momentum tensor of the source is given by

$$T^\nu_\mu = (\varepsilon + p)u_\mu u^\nu - p\delta^\nu_\mu, \quad (2.4)$$

where u^μ is the flow vector satisfying

$$g_{\mu\nu}u^\mu u^\nu = 1. \quad (2.5)$$

Here ε is the total energy density of a perfect fluid and/or dark energy density, while p is the corresponding pressure. p and ε are related by an equation of state which will be studied below in detail. In a co-moving system of coordinates from (2.4) one finds

$$T_0^0 = \varepsilon, \quad T_1^1 = T_2^2 = T_3^3 = -p. \quad (2.6)$$

In view of (2.6) from (2.3) one immediately obtains [6]

$$a(t) = D_1 \tau^{1/3} \exp\left[X_1 \int \frac{dt}{\tau(t)}\right], \quad (2.7a)$$

$$b(t) = D_2 \tau^{1/3} \exp\left[X_2 \int \frac{dt}{\tau(t)}\right], \quad (2.7b)$$

$$c(t) = D_3 \tau^{1/3} \exp\left[X_3 \int \frac{dt}{\tau(t)}\right]. \quad (2.7c)$$

Here D_i and X_i are some arbitrary constants obeying

$$D_1 D_2 D_3 = 1, \quad X_1 + X_2 + X_3 = 0,$$

and τ is a function of t defined to be

$$\tau = abc. \quad (2.8)$$

From (2.3) for τ one find

$$\frac{\ddot{\tau}}{\tau} = \frac{3\kappa}{2}(\varepsilon - p) + 3\Lambda. \quad (2.9)$$

On the other hand the conservation law for the energy-momentum tensor gives

$$\dot{\varepsilon} = -\frac{\dot{\tau}}{\tau}(\varepsilon + p). \quad (2.10)$$

After a little manipulations from (2.9) and (2.10) we find

$$\dot{\tau}^2 = 3(\kappa\varepsilon + \Lambda)\tau^2 + C_1, \quad (2.11)$$

with c_1 being an arbitrary constant. Let us now, in analogy with Hubble constant, define

$$\frac{\dot{\tau}}{\tau} = \frac{\dot{a}}{a} + \frac{\dot{b}}{b} + \frac{\dot{c}}{c} = 3H. \quad (2.12)$$

On account of (2.12) from (2.11) one derives

$$\kappa\varepsilon = 3H^2 - \Lambda - C_1/(3\tau^2). \quad (2.13)$$

It should be noted that the energy density of the Universe is a positive quantity. It is believed that at the early stage of evolution when the volume scale τ was close to zero, the energy density of the Universe was infinitely large. On the other hand with the expansion of the Universe, i.e., with the increase of τ , the energy density ε decreases and an infinitely large τ corresponds to a ε close to zero. Say at some stage of evolution ε is too small to be ignored. In that case from (2.13) follows

$$3H^2 - \Lambda \rightarrow 0. \quad (2.14)$$

As it is seen from (2.14) in this case Λ is essentially non-negative. We can also conclude from (2.14) that in absence of a Λ term beginning from some value of τ the evolution of the Universe comes stand-still, i.e., τ becomes constant, since H becomes trivial, whereas in case of a positive Λ the process of evolution of the Universe never comes to a halt. Moreover

it is believed that the presence of the dark energy (which can be explained with a positive Λ as well) results in the accelerated expansion of the Universe. As far as negative Λ is concerned, its presence imposes some restriction on ε , namely, ε can never be small enough to be ignored. It means in that case there exists some upper limit for τ as well (note that τ is essentially nonnegative, i.e. bound from below). In our previous papers we came to the same conclusion [6, 7] [with a positive Λ which in the present paper appears to be negative].

Inserting (2.12) and (2.13) into (2.9) one now finds

$$\dot{H} = -\frac{1}{2}(3H^2 - \Lambda + \frac{C_1}{3\tau^2} + \kappa p) = -\frac{\kappa}{2}(\varepsilon + p) - \frac{C_1}{3\tau^2}. \quad (2.15)$$

In view of (2.13) from (2.15) follows that if the perfect fluid is given by a stiff matter where $p = \varepsilon$, the corresponding solution does not depend on the constant C_1 .

Let us now go back to the Eq. (2.11). It is in fact the first integral of (2.9) and can be written as

$$\dot{\tau} = \pm \sqrt{C_1 + 3(\kappa\varepsilon + \Lambda)\tau^2} \quad (2.16)$$

On the other hand, rewriting (2.10) in the form

$$\frac{\dot{\varepsilon}}{\varepsilon + p} = \frac{\dot{\tau}}{\tau}, \quad (2.17)$$

and taking into account that p is a function of ε , one concludes that the right hand side of the Eq. (2.9) is a function of τ only, i.e.,

$$\ddot{\tau} = \frac{3\kappa}{2}(\varepsilon - p)\tau + 3\Lambda\tau = \mathcal{F}(\tau). \quad (2.18)$$

From a mechanical point of view Eq. (2.18) can be interpreted as an equation of motion of a single particle with unit mass under the force $\mathcal{F}(\tau)$. Then the following first integral exists [7]:

$$\dot{\tau} = \sqrt{2[\mathcal{E} - \mathcal{U}(\tau)]}. \quad (2.19)$$

Here $\mathcal{E}$ can be viewed as energy and $\mathcal{U}(\tau)$ is the potential of the force $\mathcal{F}$. Comparing the Eqs. (2.16) and (2.19) one finds $\mathcal{E} = C_1/2$ and

$$\mathcal{U}(\tau) = -\frac{3}{2}(\kappa\varepsilon + \Lambda)\tau^2. \quad (2.20)$$

Let us finally write the solution to the Eq. (2.9) in quadrature:

$$\frac{d\tau}{\sqrt{C_1 + 3(\kappa\varepsilon + \Lambda)\tau^2}} = t + t_0, \quad (2.21)$$

where the integration constant t_0 can be taken to be zero, since it only gives a shift in time.

In what follows we study the Eqs. (2.9) and (2.10) for perfect fluid obeying different equations of state.

III. UNIVERSE FILLED WITH PERFECT FLUID

In this section we consider the case when the source field is given by a perfect fluid. Here we study two possibilities: (i) the energy density and the pressure of the perfect fluid are connected by a linear equation of state; (ii) the equation of state is a nonlinear (Van der Waals) one.

A. Universe as a perfect fluid with $p_{\text{pf}} = \zeta \varepsilon_{\text{pf}}$

In this subsection we consider the case when the source field is given by a perfect fluid obeying the equation of state

$$p_{\text{pf}} = \zeta \varepsilon_{\text{pf}}, \quad (3.1)$$

where ζ is a constant and lies in the interval $\zeta \in [0, 1]$. Depending on its numerical value, ζ describes the following types of Universes [11]

$$\zeta = 0, \quad (\text{dust Universe}), \quad (3.2a)$$

$$\zeta = 1/3, \quad (\text{radiation Universe}), \quad (3.2b)$$

$$\zeta \in (1/3, 1), \quad (\text{hard Universes}), \quad (3.2c)$$

$$\zeta = 1, \quad (\text{Zel'dovich Universe or stiff matter}). \quad (3.2d)$$

In view of (3.1), from (2.10) for the energy density and pressure one obtains

$$\varepsilon_{\text{pf}} = \varepsilon_0 / \tau^{(1+\zeta)}, \quad p_{\text{pf}} = \zeta \varepsilon_0 / \tau^{(1+\zeta)}, \quad (3.3)$$

where ε_0 is the constant of integration. For τ from (2.21) one finds

$$\frac{d\tau}{\sqrt{C_1 + 3(\kappa \varepsilon_0 \tau^{1-\zeta} + \Lambda \tau^2)}} = t. \quad (3.4)$$

As one sees, the positivity of the radical in (3.4) for a negative Λ imposes some restriction on the upper value of τ , i.e., τ should be bound from above as well. In Fig. 1 the graphical view of the potential $\mathcal{U}(\tau)$ is illustrated for a negative Λ . As it was mentioned earlier, $\mathcal{E}$ or C_1 in case of $\zeta = 1$ does not play any role. Universe in this case initially expands, reaches to the maximum and then begin to contract finally giving rise to a space-time singularity [cf Fig. 2]. For the other cases depending on the choice of $\mathcal{E}$ expansion of the Universe is either non-periodic [Fig. 3] with a singularity at the end or oscillatory one without space-time singularity [Fig. 2]. In Fig. 4 we demonstrate the evolution of the BI Universe with a positive Λ . In this case the Universe expands exponentially, the initial anisotropy quickly dies away and the BI Universe evolves into a isotropic FRW one. There does not any upper bound for τ in case of a positive Λ . Note that in the Figs. (1 - 4) **d**, **r**, **h** and **s** stand for dust, radiation, hard Universe and stiff matter, respectively.

In absence of the Λ term one immediately finds

$$\tau = A t^{2/(1+\zeta)}, \quad (3.5)$$

with A being some integration constant. As one sees from (2.7), in absence of a Λ term, for $\zeta < 1$ the initially anisotropic Universe eventually evolves into an isotropic FRW one, whereas, for $\zeta = 1$, i.e., in case of stiff matter the isotropization does not take place.

B. Universe as a van der Waals fluid

Here we consider the case when the source field is given by a perfect fluid with a van der Waals equation of state in absence of dissipative process. The pressure of the van der Waals




FIG. 1: View of the potential $\mathcal{U}(\tau)$. As one sees in case of stiff matter this potential allows only non-periodic solution.




FIG. 2: Evolution of volume scale τ with a negative Λ and $C = -0.1$. As one sees, in this case the model with perfect fluid given by dust, radiation and hard Universe allow oscillation, whereas, stiff matter gives rise to a non-periodic solution.




FIG. 3: Evolution of volume scale τ with a negative Λ and $C = 0$. In this case the model with perfect fluid given by dust, radiation, hard Universe and a stiff matter gives rise to a non-periodic solution.




FIG. 4: Evolution of the Universe with a positive positive Λ . In this case independent to the choice of $zeta$ the expansion of the Universe is always exponential.

fluid p_w is related to its energy density ε_w by [1]

$$p_w = \frac{8W\varepsilon_w}{3 - \varepsilon_w} - 3\varepsilon_w^2. \quad (3.6)$$

In (3.6) the pressure and the energy density is written in terms of dimensionless reduced variables and W is a parameter connected with a reduced temperature. In the Figs. 5 and 6 the energy density and the pressure of the system are illustrated with a negative and a positive Λ term as well as in absence of it.




FIG. 5: View of energy density ε and pressure p in case of a Van der Waals fluid with a negative Λ .




FIG. 6: View of energy density ε and pressure p in case of Van der Waals fluid with $\Lambda \geq 0$.




FIG. 7: Evolution of $\tau(t)$ with the BI Universe filled with Van der Waals fluid. Independent to the sign of Λ the model provides with exponentially expanding Universe.




FIG. 8: Evolution of the BI Universe filled with different types of perfect fluid in absence of a Λ term. As one sees, in case of Van der Waals fluid $\tau(t)$ grows faster at the early stage, then slows down with time.

Inserting (3.6) into (2.15) on account of (2.13) one finds

$$\begin{aligned} \dot{H} = & -\frac{\{3H^2 - \Lambda - C_1/(3\tau^2)\}[(3 + 8W)\kappa - \{3H^2 - \Lambda - C_1/(3\tau^2)\}]}{2(3\kappa - \{3H^2 - \Lambda - C_1/(3\tau^2)\})} \\ & + \frac{3}{2\kappa}(\{3H^2 - \Lambda - C_1/(3\tau^2)\})^2. \end{aligned} \quad (3.7)$$

It can be easily verified that the Eq. (3.7) in absence of λ term and $C_1 = 0$ and $\kappa = 3$ coincides with that given in [1] :

$$\dot{H} = -\frac{3}{2}\left[H^2 + \frac{8WH^2}{3 - H^2} - 3H^4\right]. \quad (3.8)$$

The solution of the second-order differential equation (3.7) for $H(t)$ can be found by specifying the initial value for $H(t)$ at $t = 0$, for a given value of parameter W . Here we graphically present some results concerning the evolution of BI Universe with a Van der Waals fluid. In Fig. 7 we compare the evolution of τ with and without Λ term. As one sees, the character of evolution does not depend on the sign of Λ . In all cases we find exponentially expanding Universe, though the rapidity of growth depends on Λ . The Fig. 8 gives the comparison of the expansion of τ with perfect fluid obeying different equations of state.

IV. CONCLUSION

The evolution of an anisotropic Universe given by a Bianchi type I cosmological model is studied in presence of a perfect fluid and a Λ term. It has been shown that in case of a perfect fluid obeying $p = \zeta\varepsilon$, where p and ε are the pressure and energy density of the fluid, respectively, a negative Λ may generate an oscillation in the system thus giving rise to a singularity-free mode of expansion. Introduction of a positive Λ in this case results in a rapid expansion of the Universe. If the Universe is filled with a Van der Waals fluid, no oscillatory or non-periodic mode of expansion occurs. Independent to the sign of Λ the Universe in this case expands exponentially.

-
- [1] G.M. Kremer, Phys. Rev. D **68**, 123507 (2003).
 - [2] I. Zlatev, L. Wang, and P.J. Steinhardt, Phys. Rev. Lett. **85**, 896 (1999).
 - [3] J.C. Fabris, S.V.B. Goncalves, and P.E. de Souza, Gen. Relativ. Gravit. **34**, 53 (2002).
 - [4] A. Dev, J.S. Alcaniz, and D. Jain, Phys. Rev. D **67**, 023515 (2003).
 - [5] M.C. Bento, O. Bertolami, and A.a. Sen, Phys. Rev. D **66**, 043507 (2002)
 - [6] Bijan Saha, Phys. Rev. D **64**, 123501 (2001) [arXiv: gr-qc/0107013].
 - [7] Saha, Bijan and Todor Boyadjiev, Phys. Rev. D **69**, 124010 (2004) [arXiv: gr-qc/0311045].
 - [8] A. Einstein, Sitzungsber. Preuss. Acad. Wiss. **1**, 142 (1917) and **1**, 349 (1919) [English translation: *The Principle of Relativity* (Methuen, 1923, reprinted by Dover, New York, 1924), pp. 177 and 191].
 - [9] A. Guth, Physical Review D **23**, 347 (1981).
 - [10] V. Sahni, Dark Matter and Dark Energy, e-print astro-ph/0403324.
 - [11] K.C. Jacobs, Astrophysical Journal **153**, 661 (1968).